

Preparing the Infusion Bag

For Administering
RADICAVA[®] (edaravone)



Radicava[®]
(edaravone) *IV infusion*
30mg/100mL

A decorative graphic consisting of a series of colored dots (blue, green, yellow) arranged in a curved, fan-like pattern to the right of the text.

Share These Instructions


This guide shows how to properly prepare the intravenous (IV) infusion bag for administering RADICAVA[®] (edaravone). Following these instructions can help prevent damage to the stopper and avoid medication leakage.

HCPs: Please use this guide to help educate your staff and patients.

Patients: Be sure to share this guide with the care team member who is assisting with your infusions.


If you have questions about RADICAVA[®] or any medical concerns, talk to your healthcare provider.

Step 1: Align Spike



Handle the bag gently. Hold the bag with the rubber stopper facing upward. If the bag is already hanging, remove it from the IV pole. Align the spike to the center of the stopper without twisting. **DO NOT** spike upward. It is important **NOT** to insert the spike at an angle.


Step 2: Carefully Insert Spike



Very gently and carefully insert the spike straight into the center of the stopper.

It is important **NOT to twist the spike** when inserting it into the IV bag. Be sure to only spike once.

Avoid Damage to the Stopper



During insertion:

- **DO NOT** twist the spike
- **DO NOT** insert the spike at an angle
- **DO NOT** insert the spike too deep
- **DO NOT** spike multiple times

Doing so may damage the stopper and cause medication leakage.

NEVER INFUSE A BAG IF THE SPIKE HAS BEEN IMPROPERLY INSERTED INTO THE RUBBER STOPPER, CAUSING MEDICATION LEAKAGE.

For complete dosage and administration information, see full Prescribing Information and Patient Information for RADICAVA® (edaravone) available at www.RADICAVA.com.

To learn more, watch our Administration Video at RADICAVA.com/AdminVideo



Radicava[®]
(edaravone) IV infusion
30mg/100mL

Radicava[®]
(edaravone) *IV infusion*
30mg/100mL



To report product complaints, contact
Mitsubishi Tanabe Pharma America, Inc.
at **1-888-292-0058**.

RADICAVA, the RADICAVA logo, and the corporate symbol of
Mitsubishi Tanabe Pharma America are registered trademarks
of Mitsubishi Tanabe Pharma Corporation.
For US audiences only.

© 2020 Mitsubishi Tanabe Pharma America, Inc.
All rights reserved. CP-RC-US-0465 02/20



Mitsubishi Tanabe Pharma America